

Summer Work: 2020-2021

Welcome to AP Physics 1! In this course you will find out how things work and why things work the way they do. Throughout the year, we'll cover some of the major areas of physics including: Kinematics, Newton's Laws of Motion, Gravitation & Circular Motion, Work, Energy & Power, Linear Momentum, Torque & Rotational Motion, Simple Harmonic Motion, Waves & Sound, and Electrostatics & Simple Electric Circuits.

In physics, we try to understand the relationships between the physical properties of objects and systems. We use mathematics to describe these relationships. So, while this course is not a math course, we do use math quite extensively. It is extremely important to have a strong background in **basic** math and science skills to be successful in physics; therefore, your summer assignment will be to review the skills necessary for understanding the course material.

Skill 1 - Solving Equations

You will frequently need to manipulate an equation to solve for an unknown. Often the "givens" in AP Physics will not be numbers; rather they will be variables (letters). It is important that you know how to solve for any variable in an equation.

Example:

$$v_f = v_i + at^2$$

Solve the above equation above for t.

Answer:

$$t = \sqrt{\frac{v_f - v_i}{a}}$$

Directions: Solve the following equations for the variables listed below:

1. Solve for V_2 . $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$

4. Solve for R_{eq} . $\frac{1}{R_{eq}} = \frac{1}{R_1} + \frac{1}{R_2}$

2. Solve for a . $y = V_0 t + \frac{1}{2} a t^2$

5. Solve for r . $F = k \frac{Q_A Q_B}{r^2}$

Skill 2 - Describing Relationships

It is important that you understand what an equation physically means in this course. Therefore, you must be able to describe the relationships between given variables in a formula.

Example:

$$a_c = \frac{v^2}{r}$$

In the above equation, a_c equals the centripetal acceleration of an object moving in a circle with radius of r and moving around a circle with a tangential velocity of v .

- (a) In the equation above, what is the relationship between centripetal acceleration and tangential velocity?

Answer: The centripetal acceleration of the object is directly proportional to the square of the object's tangential velocity. This means that the centripetal acceleration will increase as the tangential velocity increases.

- (b) In the equation above, what is the relationship between centripetal acceleration and the radius of the circular path in which the object is traveling?

Answer: The centripetal acceleration of the object is inversely proportional to the radius of the circular path in which the object is traveling. This means that the centripetal acceleration will increase as the tangential velocity decreases.

Use the following for questions 7 – 9:

Use the following for questions 7 – 9:

In the equation at the right, F equals the electric force between two charges placed a distance r apart. Q_A is the magnitude of the first charge, Q_B is the magnitude of the second charge and k is Coulomb's constant.

$$F = k \frac{Q_A Q_B}{r^2}$$

7. In the equation above, what is the relationship between the electric force and Coulomb's constant?
8. In the equation above, what is the relationship between the electric force and the distance between the charges?
9. In the equation above, what is the relationship between the electric force and the magnitude of the first charge?

Skill 3 - Dimensional Analysis

It is important that you understand how to convert from one unit to another using conversion factors. You must know metric prefixes in order to do this. I have provided reference sheets in your summer work folder in case you have forgotten these.

Example:

Ex. How many centimeters are in 0.098 kilometers?

Answer:

$$100 \text{ cm} = 1 \text{ m}$$

$$1 \text{ km} = 1000 \text{ m}$$

$$\frac{0.098\cancel{\text{km}}}{1} \times \frac{1000\cancel{\text{m}}}{1\cancel{\text{km}}} \times \frac{100\text{cm}}{1\cancel{\text{m}}} = 9800\text{cm}$$

10. How many seconds are in 28 hours?

11. How many kiloliters are in 12,500 mL?

12. Convert 45 km/hr to m/s.

Skill 4 - Scientific Notation Scientific Notation –

Part A: There are things in physics that are very, very large (like the mass of a planet in kilograms, for example) or very, very small (like the mass of an electron in kilograms, for example). You must be able to recognize that a number is in scientific notation and know how to deal with it.

13. The following numbers are in scientific notation.
Express them in standard notation.

A. 6.370×10^4 mg

B. 4.2×10^{-2} m

14. The following numbers are in standard notation.
Express them in proper scientific notation.

A. 0.0000015 g

B. 763,420,000 cm

Scientific Notation - Part B: You will be required to use scientific notation in calculations. *Hint: I am sure you have learned the "tricks" for multiplying and dividing exponents in math class... You may use those "tricks" here.*

15. Add or subtract as indicated.

A. 3.2×10^5 cm + 4.8×10^5 cm

B. 3.29×10^5 g - 2.8×10^4 g

16. Multiply or divide as indicated.

A. $(4.0 \times 10^{-8}$ cm) x $(2.0 \times 10^5$ cm)

B. $(9.0 \times 10^7$ m) / $(3.0 \times 10^3$ s)

Skill 5 - Significant Figures

Part A: You must know what significant figures are and how to determine the number of significant figures in a measurement. I have included the rules for determining the number of sig figs in a measurement below in case you have not encountered this in any of your math or science classes yet.

What are significant figures?

In scientific work, all numbers are assumed to be derived from measurements and, therefore, the last digit in each number is uncertain. All certain digits plus the first uncertain digit are significant figures. Only numbers determined by definition or by counting are exact. Numbers determined by definition or counting are said to have an infinite number of significant figures.

Four Rules for Determining the Number of Sig Figs in a Measurement:

1. **Nonzero digits are always significant.** (Ex. There are 3 sig figs in 568 cm and 2 sig figs in 1.4 seconds.)
2. **All final zeros after a decimal point are significant.** (Ex. There are 4 sig figs in 2.300 sec.)
3. **Zeros between two other significant digits are always significant.** (Ex. There are 3 sig figs in 203 m/s and 4 sig figs in 2.002 cm.)
4. **Zeros solely used as placeholders are NOT significant.** (Ex. There are 2 sig figs in 26,000 grams and only 1 sig fig in 0.000005 km)

State the number of significant figures in each measurement:

17. 1405 kg _____

18. 0.0034 m _____

19. 5.80×10^6 kg _____

Significant Figures - Part B: You must know how to determine the number of significant figures that should be in your answer.

Rules for Determining the Number of Sig Figs in an Answer When Adding and/or Subtracting:

1. Determine the **precision** of each measurement. (Precision means how many places after the decimal for each measurement. Ex. 1.0 = one decimal place, 1.12 = two decimal places)
2. Make a note of the lowest number of decimal places. This is the least precise measurement.
3. Now add or subtract the measurements.
4. Round your answer so that it matches the precision of the measurement with the lowest number of decimal places.

Ex.

Add $1.02 \text{ s} + 0.0003 \text{ s} + 26.022 \text{ s} = ?$

1.02 has 2 decimal places

0.0003 has 4 decimal places

26.022 has 3 decimal places

So, 1.02 s is the least precise with only two decimal places - our answer must match this.

Add to find the answer:

$1.02 \text{ s} + 0.0003 \text{ s} + 26.022 \text{ s} = 27.0423 \text{ s}$

Round to two decimal places:

So, our answer with the correct amount of sig figs is

27.04 s

Rules for Determining the Number of Sig Figs in an Answer When Multiplying and/or Dividing:

1. Determine the **# of sig figs** in each measurement. (Use the rules above.)
2. Make a note of the lowest number of sig figs.
3. Now multiply or divide the measurements.
4. Round your answer so that it has the same number of sig figs as the measurement with the least amount.

Ex.

Multiply $1.02 \text{ m} \times 0.0003 \text{ m} \times 26.022 \text{ m} = ?$

1.02 has 3 sig figs

0.0003 has 1 sig fig

26.022 has 5 sig figs

Note the lowest amount of sig figs: 1 sig fig

Multiply to find the answer:

$1.02 \text{ m} \times 0.0003 \text{ m} \times 26.022 \text{ m} = 0.007962732 \text{ m}^3$

Round so that the answer only has 1 sig fig:

So, our answer with the correct amount of sig figs is

0.008 m³

Perform the following operations. Use the correct number of significant figures in your answers.

20. $(9.2 \text{ cm}) + (0.0080 \text{ cm}) + (8.30 \text{ cm})$


21. $60.000 \text{ g} \div (2.000 \text{ m} \times 3.00 \text{ m} \times 1.000 \text{ m})$

Skill 6 – Trigonometry


Not everything in physics is simply straight up and down or left and right. In the real world, we need to deal with angles. This is where trigonometry and the Pythagorean Theorem come in handy. I have the basics below in case you have not done this in your math classes yet.

There are 3 trig functions that you will use on a regular basis in physics problems: **sine**, **cosine** and **tangent**. An easy way to remember them is by using "SOH CAH TOA" Note: (See below) the opposite side is the side opposite the reference angle θ ; the hypotenuse is the longest side opposite the right angle. The adjacent side is the one that makes up part of the reference angle θ and is not the hypotenuse.


SOH $\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$
CAH $\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$
TOA $\tan \theta = \frac{\text{opposite}}{\text{adjacent}}$



The **Pythagorean Theorem** is another formula that you will use frequently in physics. $a^2 + b^2 = c^2$



22. For the triangle below, find:



A. $\sin \theta =$

B. $\cos \theta =$


C. $\tan \theta =$

Use the following trig table to answer questions 23+24. Do not use a calculator.


Ex. from table: $\sin(90) = 1$

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined


23. Calculate the value of y in the triangle below.



24. Calculate the value of x in the triangle below.




25. Calculate the value of F in the triangle below.



Skill 7 – Reading Graphs

You must be able to pull out and explain information about data from an experiment just by looking at a graph of this data. This will include recognizing relationships and trends in the data.

Graph 1: Distance a Person Walked vs. Time



Answer the following questions using the graph above.

1. What is the independent variable for this graph? What units are used to measure this quantity? _____
2. What is the dependent variable for this graph? What units are used to measure this quantity? _____
3. At what position did the person start? _____
4. What was the position of the person at 2 hours? _____
5. What was the position of the person at 4.5 hours? _____
6. Approximately, at what time (in seconds) did the person reach 4,000 meters? _____
7. Approximately, at what time (in seconds) did the person reach 14,000 meters? _____
8. At what time, in seconds, did the person reach its maximum position? _____
9. Describe the motion of the person from point P to point R. How long (in hours) did they do this for?
10. Rank the **speed** of the person during the following intervals (1 = least, 3 = greatest). Explain how you determined this.
 _____ from X to P _____ from P to R _____ from R to S

Skill 8 – Constructing Graphs (Graphing Data)

Graphs are useful tools in physics because trends in data and relationships between variables are easy to visualize when represented graphically. Now, instead of just reading a graph, you must also create your own before analyzing the data. Review the steps below for making graphs in science to be sure you have included all of the necessary components.


Following the steps below will help you to make certain that all components of the graph are correctly presented.

1. **Identify the variables.** Independent on the x-axis and dependent on the y-axis
2. **Determine the range.** What is the highest value data point for each axis?
3. **Select the scale units.** Divide each axis uniformly into appropriate units using the maximum amount of space available. (Remember that the axes may be divided differently but each square along the same axis must represent the same interval.)
4. **Number and label each axis.** Be sure to include units where appropriate as part of the axis label.
5. **Plot the data points** as ordered pairs. (x,y)
6. **Draw the best fit line (may be a straight line or a smooth curve).** For a straight line, eyeball it. "Eyeball it" means: use a straight edge to draw your line in such a way that approximately the same number of points lie above and below the line. Note: Lines in physics are not always straight! See what shape works the best from your "Relationships on Graphs - Cheat Sheet" and sketch a smooth curve to fit the data.
7. **Title the graph.** The title should clearly describe the information contained in the graph. It is common to mention the dependent variable (y-axis) first followed by the independent variable (x-axis). See title in graph on previous page.

Sample Data Set A:

The following set of data was collected while experimenting with position and time of a miniature motorized car traveling on a straight track. Plot the data below on the grid. After plotting the graph for the data set, use the graph to answer the analysis questions in Skill 3.


Time (minutes)	Position (meters)
0	0
5	18
10	32
15	45
20	58
25	74



Sample Data Set B:

The following set of data was collected during an experiment to find the density for an unknown pure metal. Five different volumes of the same unknown pure metal were massed and the data was recorded below. After plotting the graph for the data set, use the graph to answer the analysis questions in Skill 3.

Volume (cm ³)	Mass (g)
0.18	2.00
0.44	5.00
0.66	7.50
1.41	16.00
2.11	24.00



Skill 9 – Analyzing Graphs You must be able to use the graph to analyze data from an experiment. This includes predicting relationships, or using the slope, x and y-intercepts, and/or the area under the curve to tell you physical information about an experiment or relationship. This may also include interpolation or extrapolation. (P.S. Be sure you know when it is appropriate to use interpolation or extrapolation.)

Analysis Questions: Directions: Use your graph from Sample Data Set A for Questions 1-6.

1. What is the independent variable for this graph? ...the dependent variable? Explain.

2. Determine the position of the car after 2.5 minutes.

3. Is there a relationship between the variables on the x and y-axis? If so, what type of relationship is this? How do you know this?

4. Calculate the slope of the best fit line. (DO NOT USE ORIGINAL DATA POINTS UNLESS THEY LIE ON YOUR DRAWN IN BEST FIT LINE.) Show your work below.

5. The slope of the best fit line of this graph is equal to the speed of car. Explain why this is true using your graph, the definition of slope, and the definition (or formula) for speed.

6. If the experiment were carried out for 30 minutes, and the motion of the car remained as it was before, what would be the position of the car at this time?

Due: Tuesday, August 24th

Usefull links to review the material

Chapter 1 <https://flexbooks.ck12.org/cbook/ck-12-physics-flexbook-2.0/>

Chapter 1 <https://cnx.org/contents/zOZP3vRI@13.9:QP7IQWBk@7/Introduction>

Have a wonderful summer!!!